

Panoflex[®] Herbicide

with TotalSol[®] soluble granules



Enhancing winter annual control in your glyphosate or post harvest fall burndown program.

Panoflex[®] herbicide with TotalSol[®] soluble granules applied for fall burndown delivers:

- An optimal ratio of two active ingredients to maximize the burndown performance of your glyphosate program.
- Flexibility to plant corn or soybeans next spring.
- Additional burndown on glyphosate-tolerant tough winter annuals, dandelions, henbit, marehail* and chickweed.
- Easy to handle dry-flowable formulation.
- 20 oz. bottles; 60 acres at 0.33 oz./A; 33 acres at 0.6 oz./A.

MIDWEST

RATE RECOMMENDATION

Panoflex herbicide 0.6 oz./A in the fall if used alone or with glyphosate

- Adjuvants are recommended. If applied in a tank-mix combination with glyphosate that contains a built-in adjuvant system, no additional surfactant is needed.
- For additional control of grasses and broadleaves, Panoflex herbicide may be tank mixed with other burndown herbicides.

Application information

- Apply by ground or by air.
- Timing — May be applied as a burndown treatment to emerged weeds preplant after fall harvest through early spring up to planting, whenever the ground is not frozen.
- Follow FMC recommended best mixing practices for burndown applications.

* 2,4-D LVE addition required.



Panoflex® herbicide with TotalSol® soluble granules

Weeds controlled at 0.3-0.6 oz./A

Black mustard
Blue/purple mustard
Bushy wallflower/treacle mustard¹
Canada thistle²
Canola, volunteer (except Clearfield®)²
Coast fiddleneck
Common chickweed¹
Common groundsel
Common lambsquarters¹
Common purslane
Corn, gromwell²
Corn spurry
Cowcockle
Cressleaf groundsel³ (butterweed)
Curly dock²
Dandelion
Deadnettle (at 0.6 oz.)
Early whitlow grass
False chamomile/wild chamomile/scentless chamomile (*Matricaria maritima* L.)
Field pennycress
Flixweed¹
Hairy buttercup
London rocket
Marestail^{1,3}
Marshelder¹

Mayweed chamomile/stinking chamomile/
dog fennel (*Anthemis cotula* L.)^{1,2}
Miner's lettuce
Narrowleaf hawksbeard^{2,3}
Nightflowering catchfly
Pineapple weed
Poison hemlock³
Prickly lettuce^{1,2}
Puncturevine
Purslane speedwell (at 0.6 oz.)³
Redroot pigweed¹
Russian thistle^{1,2}
Shepherd's purse
Slimleaf lambsquarters
Smallflower buttercup (at 0.6 oz.)³
Small seed false flax¹
Smartweeds, annual
Tansy mustard
Tarweed fiddleneck
Tumble pigweed (at 0.6 oz.)
Tumble/Jim Hill mustard²
Velvetleaf
White cockle (at 0.6 oz.)
Wild mustard¹
Wild parsnip³

Weeds partially controlled⁴

Annual sowthistle
Common cocklebur¹
Common sunflower (volunteer)^{1,2}
Common vetch²
Eastern black nightshade¹
Hairy nightshade
Hairy vetch²
Henbit
Jimsonweed
Narrowleaf hawksbeard
Pennsylvania smartweed
Prostrate knotweed
Redmaids
Redstem filaree³
Wild buckwheat
Wild carrot
Wild garlic
Wild radish²

¹ Naturally occurring resistant biotypes are known to occur.

² See the Specific Weed Instructions section of the label for more information.

³ 2,4-D LVE addition required.

⁴ For better results, include a tank-mix partner such as 2,4-D, MCPA, bromoxynil or dicamba. See the "Tank Mixtures" section of the label.



FMC burndown program, dandelion 15 DAT.

For more information, contact your local FMC retailer or representative to learn more about Panoflex herbicide from FMC and visit us at FMCCrop.com.

Always read and follow all label directions, restrictions and precautions for use. Some products may not be registered for sale or use in all states. FMC, the FMC logo, Panoflex and TotalSol are trademarks of FMC Corporation or an affiliate. Clearfield is a trademark of BASF. ©2019 FMC Corporation. All rights reserved. 18-FMC-0982 06/19

PANOFLEX
HERBICIDE



FMC | An Agricultural
Sciences Company