

HOODED SPRAY APPLICATIONS FOR FAST ACTIVITY, MORE COMPLETE BURNDOWN AND CONTROL OF TOUGH BROADLEAF WEEDS.

BROADLEAF WEED CONTROL

Hooded or Directed Sprays	
Use Rates:	1.0 – 2.0 fl. oz./A
Tank-Mix Recommendations:	<ul style="list-style-type: none"> • Glyphosate 24 – 48 fl. oz. or Gramoxone Inteon™ 1.3 – 2.0 pts. • Residual herbicides may also be used according to their labels. • Do not drift spray mixture on desirable vegetation, green bark, vegetative parts or desirable fruit. • Target 4" or less weeds for optimum control. • Always add an adjuvant according to the adjuvant guidelines. • Use a minimum of 20 gallons finished spray per acre.

TECHNICAL INFORMATION

- Active Ingredient: Carfentrazone-ethyl




APPLICATION GUIDELINES

- 35 GPA recommended, especially when weed populations are high, or weed size is large, over 6".
- Use nozzles and spray systems designed for maximum weed coverage with low spray drift.
- Aim® EC herbicide is rainfast within one hour of application.

ADJUVANT GUIDELINES

- COC at 1% v/v is recommended under dry conditions, heavy weed pressure or MSO and silicone combinations.
- NIS may be used at 0.25% for all crops.
- Add AMS/UAN and or water conditioner.

% WEED CONTROL IN RASPBERRIES FMC IN-HOUSE DATA – ARKANSAS



% WEED CONTROL IN BLUEBERRIES DATA COURTESY DR. KATIE JENNINGS, NCSU

