Elevest[™] Insect Control

Powered by Rynaxypyr® active



Product Information:

Knock down a broad-spectrum of challenging pests with the lasting residual control of Elevest™ insect control. By maximizing the ratio of ingredients bifenthrin and Rynaxypyr® active, growers get better overall control.

This convenient, fully loaded combination is more effective on stink bugs, plant bugs and brown stink bugs than other similar premix products. It provides best-in-class Lepidopteran control, making it a great fit for growers fighting the worm complex and other insects simultaneously.

Elevest insect control offers superior rainfastness, is easy to mix and clean out of tanks. It provides excellent compatibility with commonly used tank-mix partners labeled for use on the same crops, including insecticides, fungicides, herbicides, foliar fertilizers and adjuvants.

Crops that can benefit from Elevest insect control include soybeans, especially for corn earworms, loopers and stink bug control; cotton, especially for late-season bollworm control; sweet corn, peanuts and potatoes.



Ouick Facts:

- Targets key soybean pests often found at the reproductive growth stages including armyworms, corn earworms, soybean loopers and stink bugs.
- Contains maximum active ingredient ratio of industry leading Lepidopteran active, Rynaxypyr (Group 28), with the fast knockdown of superior pyrethroid, bifenthrin (Group 3A), for broad spectrum and better overall control.
- Excellent fit for growers fighting the worm complex and other insects simultaneously.
- Provides fast knockdown of targeted pests.
- Rainfast when dry.
- Labeled for use with soybeans, cotton, sweet corn, peanuts and potatoes.
- Broad-spectrum control with lower risk for mite flares*.

*Except for locations where mite resistance to bifenthrin exists.

Target Pests:

Webworms

Armyworms Grasshoppers Plant bugs**
Bollworms Beetles Stink bugs**
Cutworms Weevils** Whiteflies**
Loopers Aphids** Thrips**
Stem and fruit borers Fleahoppers Leafminers**


Leafhoppers


Active Ingredients:

Bifenthrin, Group 3A Chlorantraniliprole, Group 28

Efficacy of Elevest™ insect control on brown stink bug (*Euschistus servus*) in soybeans.

Mites**





% Brown stink bug control across all evaluations (N=4 (Untreated = 1.81 stink bugs/25 sweeps)

Efficacy of Elevest insect control on bollworms (Helicoverpa zea) in cotton.

Mean across five trials (5-8 DAT), four trials (9-14 DAT) or three trials (15-21 DAT) conducted in 2019 in Arkansas, Louisiana, Mississippi, Georgia and North Carolina.



Source: CET-19-509, Dr. David Kerns, Texas A&M University; SOK-19-104, Dr. Dominic Reisig, NC State University; SWH-19-204, Dr. Sebe Brown, Louisiana State University; SWH-19-209, Dr. Angus Catchot, Mississippi State University; SWH-19-219, Dr. Gus Lorenz, University of Arkansas; SOK-19-106, Dr. Philip Roberts, University of Georgia

For more information about Elevest™ insect control, contact your FMC retailer or visit Ag.FMC.com.

Besiege insecticide, Brigade 2EC insecticide/miticide and Elevest insect control are Restricted Use Pesticides. Always read and follow all label directions, precautions and restrictions for use. Some products may not be registered for sale or use in all states. Elevest insect control may not be registered for sale or use in all states. Contact your local FMC retailer or representative for details and availability in your state. FMC, the FMC logo, Brigade, Elevest and Rynaxypry are trademarks of FMC Corporation or an affiliate. Besiege is a trademark of a Syngenta Group Company. Orthene is a trademark of OMS Investments, Inc. ©2020 FMC Corporation. All rights reserved. 20-FMC-0955 05/20





^{**}Some species that are normally difficult to control or may have developed some level of resistance to pyrethroids require a higher rate range of 7.7 – 9.6 fl. oz/A (0.134 – 0.167 lb. ai/A), tank mixing with another insecticide or may only be labeled at 9.6 fl. oz./A. See federal label for specific rates for each crop and pest.