

Aim® EC Herbicide – Harvest Aid – Dry Legumes Target tough weeds and immature plants for quick desiccation and easy harvest.



Aim EC Herbicide

Keep harvest running smoothly with Aim® EC herbicide. Apply Aim EC herbicide as a harvest aid in dry legumes – dry beans, dry peas, chickpeas and lentils – to desiccate or dry down stubborn weeds including kochia, lambsquarters and Russian thistle. (registered for use on all Subgroup 6C)

Proper desiccation of uneven crop maturity and weed escapes can make direct harvesting or swathing of dry legume crops a quick and easy operation.

Apply Aim EC herbicide to dry-shelled peas and beans (Crop Group 6-C) when 80-90% of the seed pods are yellow or buckskin in color and only 30% of the green leaves remain on the plant.

Quick Facts

- Zero-day PHI.
- Reliable and consistent in performance, even under adverse condition.
- Excellent tool for direct harvest dry peas, lentils and chickpeas.
- No complicated rotational restrictions.
- Fast desiccation of weed escapes, which is essential to getting your crop out of the field faster.
- Cost-effective option for optimizing your defoliation and harvest schedule.
- No export restrictions.

Aim EC herbicide improves harvestability of fields with:

- Kochia
- Tansy mustard
- Lambsquarters
- Morningglories
- Field bindweed
- Russian thistle
- Pigweeds/waterhemp
- Field pennycress

Inside this issue

July 31, 2020

Aim EC Herbicide
Harvest Aid

Weeds Controlled

Use Guidelines

Jeff Wynn
S. Idaho, Nevada, Utah
208-705-3999
Jeffery.Wynn@fmc.com

Katie Kratky
Montana and Wyoming
406-320-2239
kathryn.kratky@fmc.com

Carmine Tyler
N. Idaho, E. Washington
and E. Oregon
208-596-0206
carmine.tyler@fmc.com

Kris Thomas
Columbia Basin
509-906-0375
Kris.Thomas@fmc.com

Authored by
Kirk A. Sager
509-770-0302
kirk.sager@fmc.com
Technical Service Manager

Aim® EC Herbicide Desiccation Recommendations

Desiccation Use Rates

Use Rates: 1.0-6.1 fl. oz./A

Aim® EC herbicide may be applied alone in single application or in sequential application programs on dry peas, chickpeas and lentils.

Do not apply more than 6.1 fl. oz./A total of Aim EC herbicide per season

Recommended Desiccation Programs

Option 1 (Stand Alone): Aim EC herbicide 2.0 – 6.0 fl. oz plus adjuvant

Option 2 (Tank Mix): Aim EC herbicide 1.5 – 2.0 fl. oz./A + Gramoxone Inteon® herbicide or glyphosate plus adjuvant

Adjuvant Recommendations

- Add AMS at a minimum of 17 lbs./100
- MSO is the recommended adjuvant at 1-2 gal./100 gal. of spray solution.

Application Guidelines

- Good coverage is essential for desiccation.
- **20 GPA minimum by ground application.**
- A second application may be required in seven to 10 days.
- Use flat fan nozzles within recommended pressure ranges for best performance.
- Do not use FloodJet® nozzles or tips larger than 8008 that create a coarse spray.



Pre application



Aim EC herbicide 1.5 fl. oz. +
Paraquat 16 fl. oz. + MSO 32 fl. oz. – **4 DAT**