

Aim® EC Herbicide – Harvest Aid – Wheat

Target tough weeds and immature plants for quick desiccation and easy harvest



- Apply Aim® EC herbicide at 1-2 fl. oz./A as a tank-mix partner with 2,4-D, Roundup®, or dicamba herbicides.
- **Apply once crop is mature and grain has begun to dry down. Seven-day PHI when used alone.**
- **Thorough coverage is essential for effective harvest aid.**
- Spray volumes:
 - 15 gallons/acre ground
 - 5 gallons/acre by air is recommended.
- Fast desiccation of vines and larger broadleaf weeds

Inside this issue

July 31, 2020

Aim EC Herbicide
Harvest Aid

Weeds Controlled

Use Guidelines

Aim EC herbicide improves harvestability of fields with:

- | | |
|------------------|----------------------|
| • Kochia | • Field bindweed |
| • Tansy mustard | • Russian thistle |
| • Lambsquarters | • Pigweeds/waterhemp |
| • Morningglories | • Field pennycress |



Jeff Wynn
S. Idaho, Nevada, Utah
208-705-3999
Jeffery.Wynn@fmc.com

Katie Kratky
Montana and Wyoming
406-320-2239
kathryn.kratky@fmc.com

Carmine Tyler
N. Idaho, E. Washington
and E. Oregon
208-596-0206
carmine.tyler@fmc.com

Kris Thomas
Columbia Basin
509-906-0375
Kris.Thomas@fmc.com

Authored by
Kirk A. Sager
509-770-0302
kirk.sager@fmc.com
Technical Service Manager

Use Guidelines:

- Use higher rates and water gallonage for large weeds, especially kochia or lambsquarters
- Vines will normally desiccate within a few days; other weeds may take up to a week
- AMS at 2-4 lbs. per acre may be added
- Add a COC or MSO at 2% v/v
- Observe directions/precautions on tank-mix product labels.