

Aim® EC Herbicide

Powerful defoliant and harvest aid

COTTON

PRODUCT INFORMATION:

Aim® EC herbicide from FMC is your secret weapon come harvest. It makes cotton defoliation fast, economical and easy. Aim EC herbicide is unsurpassed for cotton defoliation, juvenile regrowth removal and weed desiccation in spindle and stripper cotton.

Aim EC herbicide also delivers fast-acting, non-residual control of the toughest broadleaf weeds, including kochia, waterhemp, lambsquarters, Russian thistle and more. Flexible and easily applied alone or with a tank-mix partner for harvest aid and defoliation.

QUICK FACTS:

- Labeled for cotton, row crops and many specialty crops.
- Can be tank mixed with other cotton harvest aid products.
- Low use rate for cotton.
Up to 1.6 fl. oz./A
- Controls and suppresses broadleaf weeds.
- Flexibility to fit into many different defoliation programs.
- Effective in desiccating weeds present in cotton crop.



Cotton Harvest Aid/Defoliant

BRAND/CHEMISTRY	REMOVE MATURE FOLIAGE	REMOVE IMMATURE FOLIAGE	REGROWTH SUPPRESSION	JUVENILE REGROWTH REMOVAL	OPEN BOLLS	DESICCATE VINES AND WEEDS
Aim® EC herbicide + adjuvant	G	E	F	E	F	G
Aim EC herbicide + thidiazuron	E	E	E	E	F	G
Aim EC herbicide + ethephon	E	E	F	E	E	G
Aim EC herbicide + thidiazuron + ethephon	E	E	E	E	E	G
Aim EC herbicide + paraquat	E	E	F	E	F	E

= Excellent
 = Good
 = Fair

Target Weeds:

Cocklebur
 Hemp sesbania
 Kochia
 Lambsquarters
 Morningglory
 Nightshade
 Palmer amaranth
 Pigweed
 Russian thistle
 Velvetleaf
 Waterhemp
 Wild buckwheat*

*According to the label for control of wild buckwheat, a tank mix such as glyphosate, glufosinate, or paraquat is required.

Application Timing:

Harvest Aid Application: Apply Aim EC herbicide alone or as a tank mixture with other cotton harvest aids to defoliate and desiccate cotton and troublesome weeds that may be present at harvest. Use a good quality spray adjuvant at the recommended rates.

Application Guidelines:

Pre-Harvest Interval: 7 days

Product Use Rate: 0.25 – 1.6. oz./A

Apply when 60-70% of bolls are open or according to the state agricultural extension service guidelines in the use area.

Restrictions:

Do not apply when conditions favor drift or when wind is above 10 mph. Do not apply more than 7.9 fl. oz./A total for preplant, in-season weed control and harvest aid.

Please see product label for a complete list of restrictions.

Do not apply within 7 days of harvest.

Re-entry interval (REI): 12 hours

Label:

Active Ingredient: Carfentrazone-ethyl (Group 14), Group E herbicide

Mode of Action: The disruption of membranes by inhibiting the action of protoporphyrinogen oxidase, causing cell death.

To learn more about Aim EC herbicide, contact your FMC retailer or visit [Ag.FMC.com](https://www.Ag.FMC.com).

Always read and follow all label directions, precautions and restrictions for use. Some products may not be registered for sale or use in all states. FMC, the FMC logo and Aim are trademarks of FMC Corporation or an affiliate. ©2023 FMC Corporation. All rights reserved. 23-FMC-1504 08/23

