
Aim® EC Herbicide



PRODUCT INFORMATION:

Up your crop protection game with Aim® EC herbicide from FMC. Labeled for corn and soybeans, Aim EC preplant burndown and preharvest herbicide delivers powerful control of the toughest broadleaf weeds. Aim EC herbicide also delivers superior no-till burndown when tank mixed with a suitable burndown partner.

Aim EC herbicide is formulated for a low use rate and is easily tank mixed with other herbicides and insecticides. With preplant burndown and harvest aid options, it provides the flexibility you need to control tough broadleaves like kochia, waterhemp, lambsquarters, Russian thistle and others.

Choose Aim EC herbicide to protect your corn and soybean fields from the toughest weeds.

OUICK FACTS:

- Labeled for both corn and soybeans.
- Preplant burndown and harvest aid application options for corn and soybeans.
- Low use rate:

Corn: Up to 2.0 fl. oz./A

Soybeans: Up to 1.5 fl. oz./A

- · Controls and suppresses key broadleaf weeds.
- Superior no-till burndown.



Target Weeds:

Cocklebur

Hemp sesbania

Kochia*

Lambsquarters

Morningglory

Nightshade

Palmer amaranth

Pigweed

Russian thistle

Velvetleaf

Waterhemp

Wild buckwheat**

Application Timing:

Apply Aim® EC herbicide alone or with other herbicides or liquid fertilizers as a burndown treatment to control or suppress weeds. For optimum performance, make applications to actively growing weeds up to 4 inches high or rosettes less than 3 inches across.

Pre-harvest Intervals:

Corn: 3 days
Soybeans: 3 days
Product Use Rate:

Corn: Up to 2.0 fl. oz./A Soybeans: Up to 1.5 fl. oz./A

Restrictions:

Corn: Do not apply more than 2.0 fl. oz., 0.031 pound of active ingredient, per acre per season including all preplant, in-crop and harvest aid applications.

Soybeans: Do not apply more than 1.5 fl. oz., 0.023 pound of active ingredient, per acre per season.

Please see product label for a complete list of restrictions.

Label:

Active Ingredient: Carfentrazone-ethyl (Group 14), Group E Herbicide

Mode of Action: The disruption of membranes by inhibiting the action of protoporphyrinogen oxidase, causing cell death.

Treatment: Aim EC herbicide 1.0 oz. + 2, 4-D LV4 0.5 pt. 14 - DAT



Treatment: 2, 4-D LV4 0.5 pt. 14 DAT




Treatment: Untreated



Burndown Tank Mixture for Glyphosate Resistant Marestail

Southern Illinois University - 2015



- Authority® First DF Herbicide 4 oz./A + 2, 4-D LV4 16 oz./A + MSO 1% v/v
- Aim EC Herbicide 1 oz./A + Authority First DF Herbicide 4 oz./A + 2,4-D LV4 16 oz./A + MSO 1% v/v
 - Authority First DF Herbicide 4 oz./A + Clarity® Herbicide 8 oz./A + MSO 1% v/v
 - Aim EC Herbicide 1 oz./A + Authority First DF Herbicide 4 oz./A + Clarity Herbicide 8 oz./A + MSO 1% v/v

To learn more about Aim EC herbicide, contact your FMC retailer or visit Ag.FMC.com.





^{*}Naturally occurring resistant biotypes are known to occur.

^{**}Controlled when used with a tank-mix partner.