

A Game Changer for Your Potato Fields

POTATOES

PRODUCT INFORMATION:

Up your harvest aid game with Aim® EC herbicide from FMC. Aim EC herbicide targets leaves and stems for complete potato desiccation.

Following Aim EC herbicide's label guidelines helps improve vine kill's success rate and, in turn, tuber quality. Vine killing benefits tuber appearance, limits tuber size and improves release from the vine at harvest. Potatoes with proper skin set maintain better skin color, lose less weight in storage and are more resistant to bruising and soft rot.

Aim EC herbicide also delivers reliable control of the toughest broadleaf weeds like kochia, lambsquarters, nightshade and more.


QUICK FACTS:

- Reliable and consistent performance, even under adverse conditions.
- Controlled, rapid defoliation of leaves and quick stem kill with no stem end browning.
- Better skin set, essential for bruise management.
- Increased stolon detachment.
- Reliable control of tuber size.
- May be used on all varieties of potatoes, including potatoes grown for seed.





**Aim EC Herbicide Desiccation
Potato Yield: Hundred Weight/Acre
Miller Research — Rupert, Idaho, 2014**



Target Weeds:

Cocklebur	Palmer amaranth
Hemp sesbania	Pigweed
Kochia	Russian thistle
Lambsquarters	Velvetleaf
Morningglory	Waterhemp
Nightshade	Wild buckwheat*

*Controlled when used with a tank-mix partner

Application Timing:

Harvest Aid Desiccation: Apply Aim® EC herbicide foliar to potatoes in the later stages of senescence for desiccation of potato foliage and vines. Aim EC herbicide will also desiccate late-season susceptible broadleaf weeds to aid tuber harvest.

Adjuvant Recommendations:

Add AMS at a minimum of 17 lbs./100 gal. MSO is the recommended adjuvant at 1-2 gal./100 gal. of spray solution. COC at 1-2 gal./100 gal. of spray solution may be used if MSO is restricted in the tank mix.

Restrictions:

Do not apply more than 11.6 fl. oz./A per crop season as a desiccant. Do not apply when conditions favor drift or wind is above 10 mph.

Label:

Active Ingredient: Carfentrazone-ethyl (Group 14), Group E herbicide

Mode of Action: The disruption of membranes by inhibiting the action of protoporphyrinogen oxidase, causing cell death.

**Aim EC Herbicide Potato Desiccation
Miller Research — Rupert, Idaho, 2014**



Untreated Check



Aim EC herbicide — 2 fl. oz./A + Reglone desiccant 16 fl. oz./A

To learn more about Aim EC herbicide, contact your FMC retailer or visit AG.FMC.COM.

Always read and follow all label directions, precautions and restrictions for use. Some products may not be registered for sale or use in all states. FMC, the FMC logo and Aim are trademarks of FMC Corporation or an affiliate. Rely is a registered trademark of BASF Corporation. Reglone is a registered trademark of a Syngenta Group company. ©2023 FMC Corporation. All rights reserved. 23-FMC-1504 06/23

