

# Express<sup>®</sup> Herbicide

with TotalSol<sup>®</sup> soluble granules



## Control tough weeds in wheat, barley, oats and triticale.

If you rotate your wheat, barley, oats and triticale, you need a herbicide that doesn't carry over. Rotation flexibility is a must. However, you also need a program that gets all the tough broadleaf weeds.

The answer: Apply a tank mix of Express<sup>®</sup> herbicide with TotalSol<sup>®</sup> soluble granules plus 2,4-D or Shark<sup>®</sup> EW herbicide\*. This tank mixture delivers the broad-spectrum control you need without giving up rotation flexibility.

### Express herbicide delivers:

- Cost-effective, broad-spectrum control of your tough weeds for a fast, easy harvest.
- Excellent crop tolerance to help maximize yields.
- Wide window of application allowing you the flexibility to work around the weather.
- Excellent rotation flexibility.
- The ability to be used in a tank mix with a variety of broadleaf herbicides such as 2,4-D, MCP or dicamba, or grass herbicides.\*\*
- Excellent activity on key problems weeds like mustards, henbit, narrowleaf hawksbeard, dandelion and many other winter annual and perennial weeds. Sharpen your tank mix performance and achieve cleaner fields with your application.\*\*

\* Shark EW herbicide tank mixes are for wheat only.

\*\* Refer to label for rotational crop restrictions. Always read and follow rotational crop restrictions of the most restrictive tank-mix partner.



TotalSol soluble granules is an advanced formulation that dissolves completely in water providing more consistent weed control with faster, easier spray tank clean-out.

### Rate and Timing Recommendations

#### For wheat, barley and triticale:

Use 0.25-0.5 oz. Express herbicide per acre. 0.5 oz./A is recommended in most situations. Two applications of Express herbicide may be made per season provided the total amount does not exceed 0.5 oz./A.

#### For oats:

Apply 0.2 oz. of Express herbicide per acre for control of light populations of labeled weeds. In oats, Express herbicide must be tank mixed with another registered herbicide.

Do not make more than one application of Express herbicide per crop season on oats.

#### For wheat, barley, oats and triticale:

Apply Express herbicide after the crop is in the two-leaf stage, but before the flag leaf is visible.

#### For spring oats:

Make applications after the crop is in the three-leaf stage, but before jointing. Do not use on Ogle, Porter or Premier varieties as crop injury can occur.

## Weeds controlled

Black mustard	Early whitlowgrass	Poison hemlock <sup>1</sup>
Blue/Purple mustard	False chamomile/Wild chamomile/Scentless chamomile ( <i>Matricaria maritima</i> L.)	Prickly lettuce <sup>2,3</sup>
Bushy wallflower (treacle mustard) <sup>2</sup>	Field pennycress	Puncturevine
Canada thistle <sup>3</sup>	Flixweed <sup>2</sup>	Purslane speedwell <sup>1</sup> (at 0.5 oz./A)
Coast fiddleneck	Hairy buttercup	Redroot pigweed <sup>2</sup>
Common Chickweed <sup>2</sup>	Kochia <sup>2,3</sup>	Russian thistle <sup>2,3</sup>
Common Groundsel	London rocket	Shepherd's purse
Common Lambsquarters <sup>2</sup>	Marestail <sup>1,2</sup>	Slimleaf lambquarters
Common Purslane	Marshelder <sup>2</sup>	Small-flower buttercup <sup>1</sup> (at 0.5 oz./A)
Corn, Gromwell <sup>3</sup>	Mayweed chamomile/stinking chamomile/dog fennel ( <i>Anthemis cotula</i> L.) <sup>2,3</sup>	Smallseed falseflax <sup>2</sup>
Corn spurry	Miners lettuce	Tansymustard
Cowcockle	Narrowleaf hawksbeard <sup>1,3</sup>	Tarweed fiddleneck
Cressleaf groundsel (butterweed) <sup>1</sup>	Nightflowering catchfly	Tumble pigweed (at 0.5 oz./A)
Curly dock <sup>3</sup>	Pineapple weed	Tumble/Jim Hill mustard <sup>3</sup>
Dandelion		White cockle (at 0.5 oz./A)
Deadnettle <sup>4</sup>		Wild mustard <sup>2</sup>

## Weeds Partially Controlled<sup>5</sup>

Annual sowthistle	Hairy nightshade	Redstem filaree <sup>1</sup>
Burning nettle <sup>3</sup>	Hairy vetch <sup>3</sup>	Wild buckwheat
Common cocklebur <sup>2</sup>	Henbit	Wild carrot
Common sunflower (volunteer) <sup>2,3</sup>	Pennsylvania smartweed	Wild garlic
Common vetch <sup>3</sup>	Prostrate knotweed	Wild radish <sup>3</sup>
Eastern black nightshade <sup>2</sup>	Redmaids	

<sup>1</sup> 2,4-D LVE addition required.

<sup>2</sup> Naturally occurring resistant biotypes are known to occur.

<sup>3</sup> See the Specific Weed Instructions section of this label for more information.

<sup>4</sup> 0.5 oz. Express herbicide only.

<sup>5</sup> Partially controlled weeds exhibit a visual reduction in numbers as well as a significant loss of vigor. For better results, use 0.375-0.5 oz. Express herbicide per acre and include a tank-mix partner such as 2,4-D, MCP, bromoxynil or dicamba. See the Tank Mixtures section of this label.

## Application method

- For ground applications, use a spray volume of at least 5 GPA with flat-fan nozzles.
- For aerial applications, use nozzle types and arrangements that provide optimum spray distribution and maximum coverage. Use 5–10 GPA in California.
- See label and consult your local retailer for further mixing and application instructions.

**For more information, contact your local FMC retailer or representative to learn more about Express herbicide from FMC and visit us at [Ag.FMC.com](http://Ag.FMC.com).**

Always read and follow all label directions, precautions and restrictions for use. Some products may not be registered for sale or use in all states. FMC, the FMC logo, Express, Shark and TotalSol are trademarks of FMC Corporation or an affiliate. ©2023 FMC Corporation. All rights reserved. 23-FMC-1504 08/23

