# **Lucento**® Fungicide



# Best-in-class protection on the key sugar beet disease while preserving yield potential.

Lucento® fungicide, powered by an exclusive combincation of active ingredients only offered by FMC, delivers premier cercospora leaf spot protection in sugar beets. Under heavy pressure, field studies demonstrate that Lucento fungicide provides excellent control of cercospora leaf spot. Lucento fungicide is easily incorporated into IPM programs and can be used in rotation with other broadspectrum fungicides as part of a resistance management plan.

## **QUICK FACTS:**

- Only premix of a proven SDHI, bixafen, and flutriafol active ingredients.
- Highly systemic fungicide with preventative control of cercospora leaf spot and powdery mildew.
- Top to bottom leaf protection and upward and outward systemic movement helps maximize disease control.
- Dual modes of action reduce risk of resistance.




### **Application Guidelines**

- Maximum rate: 5.5 fl. oz./A.
- Retreatment interval: 21 days.
- Method: Ground, air, chemigation.
- Preharvest interval (PHI): 21 days.
- Maximum applications: Two per crop year.
- Restricted entry interval (REI): 12 hours.

#### **Product Information**

**Active ingredients:** Bixafen (succinate dehydrogenase inhibitors, FRAC Group 7), flutriafol (demethylation inhibitors, FRAC Group 3)



Disease	Application
Cercospora Leaf Spot	Apply Lucento fungicide tank mixed with another broad-spectrum fungicide when disease is detected in the region or conditions are favorable for disease development. Use a rotation program of different broad-spectrum fungicides for subsequent applications. Do not use Lucento fungicide back to back or more than twice in a seasonal spray program for cercospora leaf spot.





Lucento fungicide 5.5 oz. + Topsin fungicide 10 oz. (A) fb Koverall fungicide 2 lbs. + Topguard fungicide 14 oz. (B) fb Koverall fungicide 2 lbs. (C) fb Koverall fungicide 2 lbs. + Super Tin fungicide (D)



Untreated

Super Tin and Minerva Duo fungicides are Restricited Use Pesticides. Always read and follow all label directions, precautions and restrictions for use. Some products may not be registered for sale or use in all states. FMC, the FMC logo, Koverall, Lucento and Topguard are trademarks of FMC Corporation or an affiliate. Super Tin is a trademark of United Phosphorous Inc. Topsin is a trademark of Nippon Soda Company, Ltd. Proline is a registered trademark of Bayer. Minerva is a trademark of Sipcam Agro USA, Inc. ©2023 FMC Corporation. All rights reserved. 23-FMC-1504 06/23



