

Coragen® Insect Control

Powered by Rynaxypyr® active

Verimark® Insect Control

Powered by Cyazypyr® active

Exirel® Insect Control

Powered by Cyazypyr® active

Steward® EC Insecticide




TOBACCO

Insect control for tobacco.

Verimark® insect control powered by Cyazypyr® active tray drench or Coragen® insect control powered by Rynaxypyr® active transplant water applications provide early insect plant protection from transplanting through the first 30-45 days in the field to set your crop up for success. A Coragen insect control foliar application can be used post-topping if necessary for late-season target pest control. A post-topping Coragen insect control application may be applied

late in the season for additional worm control. Exirel® insect control powered by Cyazypyr® active and Steward® EC insecticide provide two additional insect control options for worms while also adding flea beetle control. Steward EC insecticide is an alternative mode of action and makes a great rotational option for the Group 28 diamides i.e., Coragen and Exirel insect control.



Consult the label for guidance regarding best practices for insecticide resistance management.

For more information, contact your local FMC retailer or representative about insect control programs from FMC and visit us at Ag.FMC.com.

Always read and follow all label directions, precautions and restrictions for use. Some products may not be registered for sale or use in all states. FMC, the FMC logo, Coragen, Cyazypyr, Exirel, Rynaxypyr, Steward and Verimark are trademarks of FMC Corporation or an affiliate. ©2023 FMC Corporation. All rights reserved. 23-FMC-1504 09/23

