## Coragen® eVo Insect Control

Powered by Rynaxypyr<sup>®</sup> active



# Help enhance crop yield and quality with consistent, long-lasting control.

#### **Product Information:**

Coragen® eVo insect control powered by Rynaxypyr® active provides an optimized active ingredient concentration to help enhance the yield and quality of your crops by achieving consistent and long-lasting control of key pests. It stops insect feeding within minutes of ingestion and provides up to three weeks of residual control against a broad spectrum of Lepidopteran pests when used according to label rates.

#### **QUICK FACTS:**

- Long-lasting protection provides extended residual control and reduces the number of applications required throughout the season.
- Reduces pest populations and helps keep them down.
- Maximum efficiency: Higher rate range than competitive products offers maximum control of key insect pests.
- Multiple application methods including at-planting soil applications, chemigation and/or foliar applications.
- Low impact on certain beneficial insects when applied at labeled rates; excellent fit in IPM programs<sup>3</sup>.
- Rainfast once spray has dried.



### The Benefits of Using Coragen® eVo Insect Control:

- Long-lasting control of Lepidopteran pests due to an optimized formulation with nearly three times the concentration of the industry standard active ingredient Rynaxypyr® active.
- Protects tobacco, vegetables, potatoes, tomatoes, lettuce, sweet corn, alfalfa, strawberries and other crops from a broad spectrum of Lepidopteran pests plus silverleaf whitefly nymphs (suppression), leafminer larvae, Colorado potato beetles and others. Excellent control of major worm pests.
- No signal word.
- Works through ingestion or contact to deliver ovicidal<sup>1</sup>, ovi-larvicidal, larvicidal and adulticide<sup>2</sup> activity.
- Fast-acting crop protection. Insects stop feeding in minutes, helping improve crop quality and reducing routes for disease entry.

Active ingredient name: Chlorantraniliprole

Chemical class: IRAC Group 28, anthranilic diamide

Formulation: 5.0 lb. ai/gallon SC

Packaging: 1-quart container

KEY CROPS	TARGET PESTS			
Tobacco	Tobacco budworm			
Tomatoes	Beet armyworm, Leafminers (larvae) <sup>5</sup> , silverleaf whiteflie (nymphs) <sup>6</sup> , Western yellowstriped armyworm			
Potatoes	Beet armyworm, Cabbage looper			
Lettuce (Head)	Beet armyworm, Cabbage looper, Corn earworm, Diamondback moth <sup>4</sup> , Leafminers (larvae) <sup>5</sup> , Tobacco budworm, Western yellowstriped armyworm			
Alfalfa	Beet armyworm, Western yellowstriped armyworm			
Brassicas	Beet armyworm, Cabbage looper, Hawaiian beet webworm, Western yellowstriped armyworm, Diamondback moth <sup>4</sup>			

<sup>&</sup>lt;sup>4</sup> See label for additional information.

<sup>&</sup>lt;sup>6</sup> Suppression only. Use in conjunction with an effective adult whitefly control program.

TECHNICAL INFORMATION									
Active Ingredient	Group	Formulation	Use Rate	PPE	REI	Packaging	Spray Interval & PHI		
Chlorantraniliprole	Group 28 anthranilic diamide, ryanodine receptor modulator.	5 lbs./gal. suspension concentrate.	0.7 - 2.5 fl. oz/A. See label for specific rates based on crop and pests.	Applicators and other handlers must wear long-sleeved shirt and long pants, shoes plus socks.	Four hours	1-quart nonrefillable container	See label for crop- specific application and pre-harvest intervals.		

Contact your local FMC retailer or representative to learn how you can get more consistent control of key pests with Coragen eVo instect control. Visit us at Ag.FMC.com.

Always read and follow all label directions, precautions and restrictions for use. Some products may not be registered for sale or use in all states. FMC, the FMC logo, Coragen and Rynaxypyr are trademarks of FMC Corporation or an affiliate. ©2024 FMC Corporation. All rights reserved. 24-FMC-0338 02/24







<sup>&</sup>lt;sup>1</sup>Ovicidal activity on selected pest species.

<sup>&</sup>lt;sup>2</sup> Based on lab studies, adult mortality is dependent on species, application rate, exposure level and time depending on application method.

<sup>&</sup>lt;sup>3</sup> In line with integrated pest management and good agricultural practices, insecticide applications should be made when pollinators are not foraging to avoid unnecessary exposure.

<sup>&</sup>lt;sup>5</sup> Control of Liriomyza species except suppression only for L. huidobrensis and L. langei.