### **AVAUNT® EVO INSECT CONTROL**

FOR MANAGEMENT OF LYGUS IN STRAWBERRIES

IRAC Group 22A Insecticide

### **PRODUCT ATTRIBUTES**

- Unique active ingredient.
- An excellent treatment option for target pest control and preservation of beneficial insects when used in accordance with label instructions.
- Good residual control.

#### **TECHNICAL INFORMATION**

• Active Ingredient: Indoxacarb

• IRAC Group: 22A

• Formulation: Water dispersible granule

Use Rate: 6 oz./ARainfastness: 2 hrs.

• REI: 12 hrs.

#### **QUICK FACTS**

- Active on both nymph and adult lygus spp.1
- Unique MOA.
- Mode of intake ingestion/feeding.
- No activity on insects exposed to dried film.
- Binds to leaf cuticle for excellent rainfastness and residual control (5 14 days).
- Coverage is key and adjuvant is recommended.
- Excellent crop safety and tank mix compatibility.
- Won't flare mites when used as labeled.
- Compatible with biological control programs and helps conserve populations of natural enemies.

¹Suppression only

| APPLICATION GUIDELINES FOR STRAWBERRIES |  |
|---|--|
| Max Application # Per Acre Per Crop     | 4 applications per acre per crop<br>Do not apply more than 24 oz./A per calendar year  |
| Ounces Per Acre                         | 6 oz./A  |
| lbs. ai/A                               | 0.11 lbs. ai/A   |
| GPA                                     | 100 to 150 gpa   |
| Minimum Interval Between Treatments     | Minimum 7 days  An interval of 21 days or less could result in detectable residues   |
| РНІ                                     | 1 day (domestic) 3 days for Canada²  |
| Application Timing                      | Early, when lygus are first detected   |
| Adjuvants                               | Recommended for coverage   |
| DO NOT USE IF EXPORTING TO MEXICO       | FMC does not guarantee compliance with any export market. It is the sole responsibility of the user to ensure compliance with export.  Users MUST test before they ship. Many factors can impact residues (environment, application method/volume, crop stage, etc.) |


<sup>&</sup>lt;sup>2</sup>It is the responsibility of the user to ensure compliance with shipper when using Avaunt eVo insect control for export to any market.






IRAC Group 22A Insecticide

# 2023 UCCE –MARK BOLDA MANAGEMENT OF LYGUS IN STRAWBERRY – WATSONVILLE



2023 Watsonville, CA. Applications: Aug 3, Aug 10, Aug 17. All treatments included a silicone adjuvant.

# 2023 UCCE –MARK BOLDA MANAGEMENT OF LYGUS IN STRAWBERRY – WATSONVILLE



2023 Watsonville, CA. Applications: Aug 3, Aug 10, Aug 17. All treatments included a silicone adjuvant.





The EPA label for Avaunt eVo insect control contains the following statements: This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are foraging the treatment area.