

RHYME® FUNGICIDE

BROAD-SPECTRUM DISEASE CONTROL

PRODUCT ATTRIBUTES	BENEFITS
Superior Activity	Rhyme® fungicide is a highly systemic demethylation inhibitor (DMI) that gives unsurpassed disease control in tree nuts.
Excellent Control	Superior disease control. Rhyme fungicide is effective in controlling pecan scab, downy spot, anthracnose, powdery mildew and zonate leaf spot.
Crop Safety	Excellent crop safety on all varieties.
Systemic Activity	Rhyme fungicide rapidly moves through the xylem to new shoots and leaves, keeping powdery mildew and other fungal pathogens out of the canopy and protecting developing fruit.

APPLICATION GUIDELINES

- Rhyme fungicide may be tank mixed with other fungicides, insecticides and other additives unless prohibited on the tank-mix partner's label.
- Rhyme fungicide should be applied in sufficient water volume to ensure sufficient coverage.
- Use the highest labeled rate and shortest retreatment interval when disease pressure is high.
- Adjuvants may be used with Rhyme fungicide unless otherwise stated in the crop use directions.
- Rhyme fungicide is compatible with most products. However, not all have been tested.
- May be applied via air or ground.
- Apply in a minimum of 10 gallons of spray solution per acre by ground and air.
- Chemigation allowed, CA pending.



TECHNICAL INFORMATION

Active Ingredient	Flutriafol
FRAC Code & Group	FRAC Group 3; DMI (demethylation inhibitor)
Formulation	SC – Suspension Concentrate
Signal Word	Warning
Application Interval	Seven days
REI	12 hours
PHI	14 days

DISEASES AND USE RATES

Pecan Scab <i>(Fusicladium effusum)</i> Downy Spot <i>(Mycosphaerella caryigena)</i> Anthracnose <i>(Colletotrichum acutatum)</i> Powdery Mildew <i>(Microsphaera penicillata)</i> Zonate Leaf Spot <i>(Cristulariella moricola)</i>	3.5 - 7 fl. oz./A	For pecan scab control, start application at bud break. Continue application at 10-14 day pre-pollination. For all other diseases, apply preventatively or when conditions are favorable for disease development. Repeat as necessary if conditions are favorable for disease development.
---	----------------------	---

RESTRICTIONS

No single application may exceed 7 fl. oz. of product/A. Do not apply more than four applications per year. Do not apply more than 28 fl. oz. of product/A per year.