Lucento° Fungicide



Best-in-class protection on key sugar beet diseases while preserving yield potential.

Powered by a novel premix of active ingredients from FMC, Lucento® fungicide provides an exclusive combination of benefits for smart disease protection and high-yielding performance against the toughest diseases in sugar beets. Under heavy pressure, field studies demonstrated that Lucento fungicide provided excellent control of cercospora leaf spot. Lucento fungicide should always be used in combination with another broad-spectrum fungicide as part of a resistance management plan.

QUICK FACTS:

Only premix of a proven SDHI, bixafen, and FMC-patented flutriafol active ingredients.

- Highly systemic fungicide with preventative control of cercospora leaf spot and powdery mildew
- Both acropetal and translaminar movement improve coverage and disease control.
- Dual modes of action reduce risk of resistance.




Application Guidelines

- Maximum rate: 5.5 fl. oz./A.
- Retreatment interval: 21 days.
- Method: Ground, air, chemigation.
- Preharvest interval (PHI): 21 days.
- Maximum applications: Two per crop year.
- Restricted entry interval (REI): 12 hours.

Product Information

Active ingredients: Bixafen (succinate dehydrogenase inhibitors, FRAC Group 7), flutriafol (demethylation inhibitors, FRAC Group 3)



	Disease	Local Recommendations
(Apply Lucento fungicide tank mixed with another broad-spectrum fungicide when disease is detected in the region or conditions are favorable for disease development. Use a rotation program of different broad-spectrum fungicides for subsequent applications. Do not use Lucento fungicide back to back or more than twice in a seasonal spray program for cercospora leaf spot.





Lucento fungicide 5.5 oz. + Topsin fungicide 10 oz. (A) fb Koverall $^{\circ}$ fungicide 2 lbs. + Topguard fungicide 14 oz. (B) fb Koverall fungicide 2 lbs. (C) fb Koverall fungicide 2 lbs. + Super Tin fungicide (D)



Untreated

Super Tin fungicide is a Restricted Use Pesticide. Always read and follow all label directions, precautions and restrictions for use. Some products may not be registered for sale or use in all states. FMC, the FMC logo, Lucento and Topguard are trademarks of FMC Corporation or an affiliate. Priaxor and Provysol are trademarks of BASF. Super Tin, Manzate and Pro Stick are trademarks of United Phosphorous, Inc. Badge is a registered trademark of Isagro USA, Inc. Topsin is a trademark of Nippon Soda Company, Ltd. Proline is a registered trademark of Bayer. Inspire is a trademark of Syngenta Group Company. Koverall is a trademark of Indofil Industries Limited. ©2021 FMC Corporation. All rights reserved. 21-FMC-0472 02/21



